

**Abstracts of the
2017 American Dairy Science Association®
Annual Meeting**

**June 25–28, 2017
Pittsburgh, PA**

***Journal of Dairy Science*®
Volume 100, Supplement 2**

Breeding and Genetics III: Methods

458 Phenotypic analysis of daily milk, fat, and protein production with geometric morphometrics. Á. A. D. Benítez^{*1}, J. I. Weller¹, and E. Ezra², ¹*Institute of Animal Sciences, Agricultural Research Organization, The Volcani Center, Rishon LeZion, Israel*, ²*Israel Cattle Breeders Association, Caesaria Industrial Park, Israel*.

Concerns have been raised in the past about the Dairy Herd Improvement Association recording frequency, because the interval between samples, about 4 weeks, may not capture the peak production for cows with shorter lactation (less than 10 mo), which have led to the conclusion that these cows have an atypical lactation curve shape. This may be due to sampling frequency rather than biological differences of cows or the influence of biotics and abiotics variables. Geometric morphometrics is a methodology that is used to measure biological shapes and curves, which has already proved its application in biology, medicine, and engineering. This methodology can be used for quantifying, testing, and visualizing shape variation and its covariation with biotic and abiotic variables. We propose to apply this method to determine the variation and covariation of lactation curves for milk, fat, and protein production. Daily records of milk production and fat and protein concentration collected by the AfiLab recording system (AfiMilk, Kibbutz Afikim, Israel) from January 2014 to January 2017 from 47 large kibbutz (communal) herds distributed throughout Israel, will be analyzed. Lactation data will be plotted into an orbital graph to depict a closed curve. Currently methods to predict future lactation prediction of individual cows are based chiefly on production on the last available test day. Using geometric morphometrics it should be possible to derive more accurate estimates of future production, which can be used both to improve management decisions and genetic evaluations.

Key Words: geometric morphometrics, milk prediction, daily fat recording

459 Genetic parameters of bovine milk color and processing characteristics predicted by mid-infrared spectroscopy. G. Visentin^{*1,2}, D. P. Berry², M. De Marchi¹, S. McParland², A. McDermott^{1,2}, S. Scarso¹, M. A. Fenelon³, and M. Penasa¹, ¹*Department of Agronomy, Food, Natural Resources, Animals, and Environment (DAFNAE), University of Padova, Legnaro (PD), Italy*, ²*Animal and Grassland Research and Innovation Center, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland*, ³*Teagasc Food Research Center, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland*.

Milk color and processing traits are important factors informing the potential and ease to manufacture milk into different dairy products. The objective of the present study was to estimate (co)variance components of milk lightness (L^*), redness-greenness (a^*), yellowness-blueness (b^*), rennet coagulation time (RCT), curd-firming time (k_{20}), curd firmness (a_{30} and a_{60}), heat coagulation time (HCT), casein micelle size (CMS), and pH, measured by mid-infrared spectroscopy on 136,807 test-day records from 9,824 Irish dairy cows between 5 and 305 d-in-milk (DIM) from parities ≤ 10 . Cow breed was defined as the proportion of Holstein, Friesian, Jersey, Norwegian Red, Montbéliarde, and "Other." Random regression models using Legendre polynomials were performed to describe the change of both additive genetic and within-lactation permanent environmental variances across different DIM. Heritability estimates averaged across all DIM for milk color were 0.31 (L^*), 0.11 (a^*), and 0.42 (b^*); average heritability estimates for processing traits ranged from 0.31 (pH) to 0.49 (k_{20}), except for HCT (0.17). Within-

trait genetic correlations approached unity between adjacent DIM, and were <0.40 at the peripheries of lactation. Eigenvalues and associated eigenfunctions of the additive genetic variance of all traits revealed that at least 80% of the total variation was associated with the height of the lactation profile. Average genetic correlations between color traits across all DIM were the weakest between a^* and b^* (-0.19); fat concentration was strongly genetically associated with b^* (0.91), while milk yield was negatively genetically correlated with all color traits. On average, RCT was strongly genetically correlated with both a_{30} (-0.68) and pH (0.75); genetic correlations between HCT and the other processing traits were, on average, weak ranging from -0.02 (with pH) to 0.28 (with RCT). Milk yield was genetically correlated with both RCT (0.31) and a_{30} (-0.49). Breeding for milk color and processing traits is possible but with some negative impact on genetic gain for milk yield.

Key Words: milk quality, breeding, infrared spectrometry

460 Genetic parameters of milk fatty acid profile in dairy sheep. J. Serdino, F. Correddu, M. G. Manca, A. Puledda, C. Dimauro, A. Nudda, and N. P. P. Macciotta^{*}, *University of Sassari, Sassari, Italy*.

Increasing consumer concerns on dairy product nutritional quality have stressed the importance of some features of milk such as the fatty acid (FA) profile. In this study, genetic parameters of milk FA profile of dairy sheep are investigated to evaluate the feasibility of breeding for improving the nutritional quality of sheep milk. Individual milk samples of 989 Sarda ewes farmed in 47 flocks located in the 4 provinces of the Island of Sardinia, Italy, were analyzed for FA composition by gas-chromatography. Genetic parameters of 15 FA (expressed as g/100 g of FA methyl ester) were estimated with an animal model, including fixed effects of lambing type, lambing month, altitude of flock, lactation stage, province and parity, and the random effects of flock-test date (FTD), and animal additive genetic. A generally high contribution of FTD to the phenotypic variance was observed (on average 51%). Heritability (h^2) estimates ranged from 0.03 for C18:3 $n-3$ to 0.48 for C16:0. Saturated and unsaturated C18 FA showed moderate to low values of h^2 (from 0.22 to 0.03, for the same FA). On the other hand they exhibited a large contribution of FTD, ranging from 0.46 for C18:0 to 0.82 for C18:3 $n-3$, respectively. The high heritability estimate of C16:0 reflects the probable genetic control of its milk content, being partly synthesized de novo in mammary gland. Genetic correlations were negative among C4:0 and short and medium chain SFA. C16:0 showed a negative correlation with most of the investigated FA, and a positive correlation with C4:0, C14:0, C14:0c9 and C16:1c9. The high heritability of C16:0, considered harmful for human health, and its negative genetic correlations with unsaturated C18 FA could be used in genetic strategies to improve the nutritional properties of milk.

Key Words: fatty acids, genetic parameters, sheep milk

461 Genomic predictions for crossbreds from all-breed data. M. E. Tooker^{*}, P. M. VanRaden, and G. C. Fok, *Animal Genomics and Improvement Laboratory, USDA-ARS, Beltsville, MD*.

Genomic predictions of transmitting ability (GPTAs) for crossbred animals were computed from marker effects of 5 dairy breeds weighted by each breed's genomic contribution to the crossbreds. Estimates of genomic breed composition are labeled breed base representation (BBR)

and are reported since May 2016 for all 1.6 million genotyped dairy animals. Animals with > 94% of any breed were rounded to 100%, and contributions of other breeds were set to 0%. All-breed scale GPTAs were first computed for each pure breed for traits milk, fat, protein, productive life, somatic cell score, daughter pregnancy rate, cow conception rate, livability, and net merit. These estimates included foreign information from multi-trait across-country evaluation (MACE) and foreign dams converted from within-breed to the all-breed base. Then, marker effects for each breed were blended by BBR to compute evaluations for crossbreds (<94% purebred) for those same traits. Conformation traits do not have an all-breed scale, so only the Jersey marker effects were applied to the crossbreds, and results seemed reasonable. Calving traits are not predicted for crossbreds, and instead a common mean was used for all crossbreds as is the current practice for breeds other than Holstein and Brown Swiss. All-breed GPTAs were then converted to within-breed GPTAs. Correlations of GPTAs for purebreds computed on the all-breed vs. current within-breed scales were 0.97 to 0.99 for most traits and breeds. Crossbred GPTAs were then computed for 44,023 crossbreds, 20,367 of which had no previous GPTAs because of breed check edits. The new GPTAs were for 1,822 Jersey × Holstein crossbreds with >40% of both breeds (F1 crosses), 75 Brown Swiss × Holstein F1, 7,237 Holstein backcrosses with >67% and <94% Holstein, 7,820 Jersey backcrosses, 313 Brown Swiss backcrosses, 1,763 other crossbreds of various mixtures, and 1,337 purebreds that had previously failed breed checks. Additional automation and redesign of many downstream programs is required for the new all-breed system to be used in weekly, monthly, and full releases. The new system is expected to provide accurate predictions for crosses among the 5 dairy breeds evaluated.

Key Words: crossbreeding, genomic prediction, breed composition

462 Genetic trends from single-step GBLUP and traditional BLUP for production traits in US Holstein. Y. Masuda*¹, I. Misztal¹, P. M. VanRaden², and T. J. Lawlor³, ¹University of Georgia, Athens, GA, ²USDA, AGIL, Beltsville MD, ³Holstein Association USA Inc., Brattleboro, VT.

The objective of this study was to compare genetic trends from a single-step genomic BLUP (ssGBLUP) and the traditional BLUP (tradBLUP) models for milk production traits in US Holstein. We used 764,029 genotyped animals in this study. Phenotypes were 305-d milk, fat, and protein yield from 21,527,040 cows recorded between January, 1990 and August, 2015. The pedigree file included 29,651,623 animals limited to 3 generations back from recorded or genotyped animals. We applied a 3-trait repeatability model with the same genetic parameters used in the US official genetic evaluation. Unknown parent groups were incorporated into the inverse of a relationship matrix (H^{-1} in ssGBLUP and A^{-1} in tradBLUP) with the QP-transformation. In ssGBLUP, 18,359 genotyped animals were randomly chosen as core animals to calculate the inverse of genomic relationship matrix with the APY algorithm. Computations with tradBLUP took 6.5 h and 1.4 GB of memory, and computations with ssGBLUP took 13 h and 115 GB of memory. Estimated breeding values were adjusted to a genetic base on recorded cows born in 2000 in each model and converted to GPTA in ssGBLUP and PTA in tradBLUP. For genotyped sires with at least 50 daughters with phenotype(s) born between 2000 and 2010, the genetic trend of GPTA was always greater than PTA in all traits. The difference in 2 genetic trends was almost constant for the sires born up to 2008 (on average, 11 kg in milk, 0.5 kg in fat, and 0.3 kg in protein yield) and the difference was greater in the last 2 years. The difference between the GPTA means for the bulls born in 2010 was 35 kg for milk, 2.2 kg for fat, and 1.2 kg for protein yield. For genotyped cows with phenotype(s), the GPTA trend was identical

to or slightly greater than the PTA trend up to 2006. Two trends started to diverge obviously in 2007 and the GPTA trend kept rising while the PTA trend remained at the same level. The single-step method provides very similar genetic trends to the traditional evaluations except for the last few years. The recent lower PTA trend can be due to a downward bias caused with genomic pre-selection of young animals.

Key Words: genomic evaluation, genetic trend, PTA

463 A Genetic Diversity Index method to improve imputation accuracies of rare variants. A. M. Butty*¹, F. Miglior^{1,2}, P. Stothard³, F. S. Schenkel¹, B. Gredler⁴, M. Sargolzaei^{1,5}, and C. F. Baes¹, ¹Centre for Genetic Improvement of Livestock, Department of Animal Biosciences, University of Guelph, Guelph, ON, Canada, ²Canadian Dairy Network, Guelph, ON, Canada, ³Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, AB, Canada, ⁴Qualitas AG, Zug, ZG, Switzerland, ⁵Semex Alliance, Guelph, ON, Canada.

Different methods to select animals for sequencing have been developed, which rely on pedigree-based relationship matrices, genomic relationships matrices, or on haplotype frequencies. Relationship-based methods select representative key animals of a population whereas haplotype frequency methods aim for better coverage of rare variants. Good average accuracies of imputation from SNP chip to whole-genome sequence (WGS) for common haplotypes were reached with the relationship-based methods. Imputation of rare variants, however, still needs to be improved, which can possibly be accomplished with a newly developed Genetic Diversity Index (GDI). This algorithm optimizes the count of unique haplotypes present in a group of animals composed of already sequenced individuals and a fixed number of sequencing candidates. Optimization is run iteratively, exchanging one candidate at a time and computing the GDI of the new group. Use of the simulated annealing algorithm defines whether the last individual added to the group should be kept. Simulated annealing has the advantage of searching for a global optimum in a situation where multiple local optima are present. The previously mentioned key ancestor and haplotype-based methods for selecting sequencing candidate were assessed and compared with the GDI algorithm using simulated cattle WGS data. Average squared correlation coefficients were used to assess imputation accuracy. A preliminary study showed that the accuracy was 1.5% higher when using GDI to enlarge the reference population than the second-best method. Application of the different methods of selection in North American Holstein data showed that the GDI algorithm selected animals carrying a higher percentage of rare haplotypes than other methods examined. Principal component analysis of the population showed that the animals selected with all tested methods were similarly distributed over the pool of candidates. When representative animals of a population are already sequenced and good overall imputation accuracies are reached, sequencing of genetically diverse animals improved the accuracy of the imputation of rare variants to the WGS density level.

Key Words: sequencing, simulation, imputation

464 Determination of quantitative trait variants by concordance via application of the a posteriori granddaughter design to the US Holstein population. J. I. Weller*^{1,2}, D. M. Bickhart², G. R. Wiggins^{2,3}, M. E. Tooker², J. R. O'Connell⁴, J. Jiang⁵, and P. M. VanRaden², ¹Agricultural Research Organization, The Volcani Center, Rishon LeZion, Israel, ²Agricultural Research Service, Beltsville, MD, ³Council on Dairy Cattle Breeding, Bowie, MD, ⁴University of